

Short-chain fatty acid PLUS assay

Study the interaction between diet, microbiome, and energy metabolism

biocrates.com



biocrates' specialized targeted assays



Comprehensive
portfolio



Reliable &
reproducible



Project
management



Data analysis &
interpretation


For research use only | not for use in diagnostic procedures

Short-chain fatty acid PLUS (SCFA+) assay

The ideal triad for studying metabolic health and the microbiome


Short-chain fatty acids (SCFA)

Microbial metabolites involved in metabolic health, energy metabolism, and immune balance.



Branched-chain fatty acids (BCFA)

Diet-based gut bacterial mediators of the gut-brain axis.



Medium-chain fatty acids (MCFA)

Energy carriers involved in insulin-resistance and the gut-immune axis.



Covered metabolites

Analyte (short name)	LLOQ* [μ M] in plasma serum	LLOQ* [nmol/g] in feces
Acetic acid (FA 2:0)	12	180
Propionic acid (FA 3:0)	4.8	72
Isobutyric acid (FA 3:0-2M)*	0.48	7.20
Butyric acid (FA 4:0)	9.6	720
2-Methylbutyric acid (FA 4:0-2M)*	0.48	7.20
Isovaleric acid (FA 4:0-3M)*	0.48	7.20
Valeric acid (FA 5:0)	0.96	14.4
2-Methylvaleric acid (FA 5:0-2M)*	0.024	0.36
3-Methylvaleric acid (FA 5:0-3M)*	0.024	0.36
Isocaproic acid (FA 5:0-4M)*	0.024	0.36
Caproic acid (FA 6:0)	0.72	10.8
4-Methylhexanoic acid (FA 6:0-4M)*	0.024	0.36
5-Methylhexanoic acid (FA 6:0-5M)*	0.024	0.36
Heptanoic acid (FA 7:0)	0.12	1.80
Caprylic acid (FA 8:0)	0.06	0.90
Nonanoic acid (FA 9:0)	0.12	1.80
Capric acid (FA 10:0)	0.12	1.80
Undecanoic acid (FA 11:0)	0.024	0.36
Lauric acid (FA 12:0)	0.48	7.20

11 SCFAs
8 MCFAs

Quantify up to 19 metabolites and 19 metabolism indicators

Standardized, quality-controlled assay with absolute quantification, optimized sample handling and normalization protocols for different matrices.

Analytical details

- LC-MS-based assay
- Robust 7-point calibration
- 3 levels of quality controls

Sample requirements

- 60 μ L plasma/serum
- 200-500 mg feces, wet weight (minimum technical amount: 50 mg)
- 3-spot sampling (recommended): 3 x 200-500 mg feces
- For other matrices, please contact us (<https://biocrates.com/contact>)

* Branched-chain fatty acid
* Lower limit of quantification

Read more



Metabolic phenotyping services

Get more information on how to produce reproducible metabolomics data

biocrates.com/services



Applications of metabolomics

Learn how metabolomics is already changing lives

biocrates.com/applications